



Intercoastal Gastroenterology Group

2 DAY Preparation for Colonoscopy (Using Clenpiq Kit)

Place: _____
Date _____
Time _____
Arrival _____
This is subject to change.

If you start any weight loss or diabetes medications such as semaglutide (Ozempic, Wegovy, etc.), tirzepatide (Mounjaro, Zepbound) or similar – please inform us as you may need to hold these prior to the procedure.

Fill your prescription for Clenpiq at your pharmacy.
Pick up (1) 10oz bottle of Magnesium Citrate OTC

The **Surgery Center** will contact you the day prior with a confirmed arrival time.

5 Days prior to procedure- Please **STOP** Advil, Motrin and Anti-inflammatory drugs, fiber supplements, iron, fish oil, vitamin E, and multivitamins. **Tylenol** is allowed. Also avoid all nuts, seeds in or on foods including popcorn, corn, granola mix, grape nuts, grapes, and raisins.

IF YOU ARE TAKING BLOOD THINNERS OR WEIGHT LOSS MEDICATIONS PLEASE NOTIFY THE GASTROENTEROLOGIST OR NURSE FOR SPECIFIC INSTRUCTIONS. _____

2 Days prior to procedure You may have a light breakfast and light lunch **ONLY** then a clear liquid diet. AT 8:00pm drink the 10oz bottle of magnesium citrate. Continue with clear liquids.

The Day prior to procedure CLEAR LIQUID DIET ONLY.

Hydration is part of your prep! Drink 6 oz. of clear liquids every hour throughout the day before you start your prep.

CLEAR LIQUID DIET Clear liquids include coffee/tea with or without sweetener. (No MILK or MILK products or CREAMERS) You may have water and flavored-waters such as; Gatorade, Kool-Aid, soda, Jell-O, and popsicles. (Without red or purple coloring) Apple juice, white grape juice, iced tea and lemonade (Without the pulp). Chicken, beef, or vegetable broth/bouillon. **No Alcohol**

Diabetic medications can be taken in the morning only. If you take insulin, monitor blood sugars and sliding scale; take only half dose of scheduled insulin in the evening. Call your PCP or Endocrinologist if questions regarding your insulin.

AT 6pm

1. Drink (1) bottle of Clenpiq.
2. Followed by (5) 8oz glass of juice or water. (finish liquids over the next 5 hours)

Continue to drink plenty of water and clear liquids through the rest of the evening.



PLEASE REMEMBER TO SET YOUR ALARM CLOCK

AT _____ 6 hours prior to procedure time, drink second bottle of Clenpiq, followed by (3) 8oz glasses of juice or water (finish liquid over the next 2 hours) **Nothing by mouth 4 hours prior to procedure.**

Morning of procedure Take all **blood pressure** and/or **cardiac medications** that you may normally take in the morning with a sip of water. (Excluding diuretics) Seizure meds and anxiety meds may also be taken.

You **MAY NOT DRIVE YOURSELF HOME** or **TAKE A TAXI** from procedure. Please make arrangements with a family member or friend. If you have any questions concerning prep or procedure, or need to reschedule, please call 342-8892. _____

YOU ARE RESPONSIBLE FOR CHECKING WITH YOUR INSURANCE COMPANY REGARDING COVERAGE FOR YOUR SURGICAL PROCEDURE. ANY DEDUCTIBLE OR COPAY IS APPLICABLE IS DUE NO LATER THAN 5 BUSINESS DAYS PRIOR TO THE DATE OF SURGERY.

IN ADDITION TO PHYSICIAN FEE, THERE IS THE FACILITY, ANESTHESIA AND POSSIBLE PATHOLOGY FEES.